ELSEVIER

Colloids and Surfaces
A: Physicochemical and Engineering Aspects 183–185 (2001) 831–834

COLLOIDS AND SURFACES A

www.elsevier.nl/locate/colsurfa

Contents of Volumes 183–185

Vols. 183–185

Editorial 1
Linear viscoelasticity of salt-free wormlike micellar solutions
S.J. Candau (Strasbourg, France) and R. Oda (Talence, France) 5
Adsorption of ammonium perfluorooctanoate at the air–water interface
J.R. Lu (Guildford, UK), R.H. Ottewill (Bristol, UK) and A.R. Rennie (London, UK) 15
Polyelectrolyte multilayer capsules templated on biological cells: core oxidation influences layer chemistry
S. Moya, L. Dähne, A. Voigt, S. Leporatti, E. Donath (Berlin, Germany) and H. Möhwald (Golm/Potsdam, Germany) 27
Interaction of poloxamer block copolymers with cosolvents and surfactants
R. Ivanova (Lund, Sweden; Ilmenau, Germany), P. Alexandridis (Lund, Sweden; Buffalo, NY, USA) and B. Lindman (Lund, Sweden) 41
Evidence of micelle growth in aqueous solutions of the amphiphilic poly(ethylene oxide)–poly(propylene oxide)–poly(ethylene oxide) triblock copolymers from differential scanning microcalorimetry
B. Michels, G. Waton and R. Zana (Strasbourg, France) 55
Microscopic study of chiral interactions in langmuir monolayer: monolayers of *N*-palmitoyl aspartic acid and *N*-stearoyl serine methyl ester
N. Nandi and D. Vollhardt (Potsdam, Germany) 67
A laser interferometer for the study of electrically excited bubble capillary modes
T. Bellini (Milano, Pavia, Italy), M. Buscaglia (Pavia, Italy), M. Corti (Segrate, Pavia, Italy), F. Mantegazza (Monza, Pavia, Italy) and C. Rocca (Pavia, Italy) 85
Interaction, critical, percolation and kinetic glass transitions in pluronic L-64 micellar solutions
S.-H. Chen, C. Liao, E. Fratini, P. Baglioni and F. Mallamace (Cambridge, MA, USA) 95
Digital cryogenic transmission electron microscopy: an advanced tool for direct imaging of complex fluids
D. Danino, A. Bernheim-Groswasser and Y. Talmon (Haifa, Israel) 113
Shear induced deformation of microcapsules: shape oscillations and membrane folding
A. Walter, H. Rehage and H. Leonhard (Essen, Germany) 123
Dynamical properties in dense triblock copolymer micellar system
P. Gambadauro, V. Venuti (Messina, Italy), F. Mallamace, C. Liao (Cambridge, MA, USA), P. Tartaglia (Cambridge, MA, USA; Roma, Italy) and S.-H. Chen (Cambridge, MA, USA) 133
Shear stimulated adsorption of surfactants from micellar solutions
A. Bakx, A.M.D.E. Timmerman and G. Frens (Delft, The Netherlands) 149
Shape fluctuations of microemulsion droplets: a neutron spin–echo study
T. Hellweg (Chemnitz, Germany), M. Gradzielski (Bayreuth, Germany), B. Farago (Grenoble, France) and D. Langevin (Orsay, France) 159
Salt-induced phase separation in the liquid crystalline phase of phosphatidylcholines
M. Rappolt, G. Pabst, H. Amenitsch and P. Laggner (Graz, Austria) 171
The electric birefringence of polyelectrolytes. Maxwell–Wagner and electrokinetic relaxation
V. Degiorgio (Pavia, Italy), T. Bellini (Milan, Italy) and F. Mantegazza (Monza, Italy) 183
Dissolution of nonionic surfactant mixtures
B.-H. Chen, C.A. Miller (Houston, TX, USA) and P.R. Garrett (Bebington, UK) 191
Stress relaxation in model transient networks: percolation and rearrangement of the crosslinks
M. Filali (Fes, Morocco), E. Michel, S. Mora, F. Molino and G. Porte (Montpellier, France) 203
Phase behavior of iridescent phases with calcium dodecyl sulfate, octanol or decanol, and water, and the effect of the presence of sodium ions
A. Zapf, R. Hammel and G. Platz (Bayreuth, Germany) 213
Induction of a lamellar phase by the addition of a hydrocarbon to an isotropic ternary surfactant system
A. Hufnagl and M. Gradzielski (Bayreuth, Germany) 227
1-Butanol and 3,3-dimethyl-1-butanol as cosurfactants of the laurylsulfobetaine/water system
M. Valiente and M. Álvarez (Madrid, Spain) 235

Oxidation of azo dyes in oil-in-water microemulsions catalyzed by metalloporphyrins in presence of lipophilic acids
M. Häger (Stockholm, Sweden), K. Holmberg (Göteborg, Sweden), A.M.A. Rocha Gonsalves (Coimbra, Portugal) and A.C. Serra (Figueira da Foz, Portugal) 247
Kinetics of surfactant adsorption: the free energy approach
H. Diamant, G. Ariel and D. Andelman (Tel Aviv, Israel) 259
PEO-related block copolymer surfactants
K. Mortensen (Roskilde, Denmark) 277
Swelling and dissolution of onion phases: the effect of temperature
M. Buchanan, S.U. Egelhaaf and M.E. Cates (Edinburgh, UK) 293
Dynamic light scattering study of polystyrene latex suspended in water at high temperatures and high pressures
R.G. Alargova, S. Deguchi and K. Tsujii (Yokosuka, Japan) 303
The gelation mechanism of solutions of isotactic polystyrene in *cis*-decalin
J. De Rudder, H. Berghmans, T. Roels and A. Jacobs (Heverlee, Belgium) 313
The vibrational motions of particle gels
K. Dawson, G. Foffi and E. Zaccarelli (Dublin, Ireland) 327
High frequency elastic modulus of hairy particle dispersions in relation to their microstructure
M.H.G. Duits, P.A. Nommensen, D. van den Ende and J. Mellema (Enschede, The Netherlands) 335
Entropy and droplet size distributions of Winsor I and II microemulsions
J.C. Eriksson, S. Ljunggren (Stockholm, Sweden), W.K. Kegel and H.N.W. Lekkerkerker (Utrecht, Netherlands) . . . 347
Small-angle neutron scattering from tetradecyltrimethylammonium bromide in NaBr aqueous solutions
G. Eckold (Gottingen, Germany) and N. Gorski (Gottingen, Germany; Moscow Region, Russian Federation) 361
Formation of cubic-phase microemulsions in sucrose alkanoate systems
K. Aramaki, M.H. Kabir, N. Nakamura, H. Kunieda and M. Ishitobi (Yokohama, Japan) 371
Kinetics of adsorption of globular proteins at liquid/fluid interfaces
R. Miller (Golm, Germany), E.V. Aksenenko (Kiev, Ukraine), V.B. Fainerman (Donetsk, Ukraine) and U. Pison (Berlin, Germany) 381
Chemically modified lipids — a suitable tool to study molecular interactions in model systems
F. Bringezu (Santa Barbara, CA, USA) and G. Brezesinski (Golm/Potsdam, Germany) 391
Miscibility of binary surfactant mixtures in the adsorbed film and micelle: the effect of size of polar head group
A. Ohta (Ishikawa, Japan), H. Matsubara, N. Ikeda and M. Aratono (Fukuoka, Japan) 403
Solubilization of benzene derivatives in silicate-surfactant systems
A. Lind, J. Andersson, S. Karlsson, M. Lindén and J.B. Rosenholm (Turku, Finland) 415
Nonconventional first order transitions in insoluble monolayers of surfactants
E. Ruckenstein (Buffalo, NY, USA) 423
Interactions between cationically modified hydroxyethyl cellulose and oppositely charged surfactants studied by gel swelling experiments — effects of surfactant type, hydrophobic modification and added salt
J. Sjöström and L. Piculell (Lund, Sweden) 429
Phase behaviors and light scattering for dilute mixtures of $C_{12}E_5$ and lauric acid
N. Awata, K. Minewaki, K. Fukada, M. Fujii and T. Kato (Tokyo, Japan) 449
Influences of monomer and oligomer structures on their adsorption on metal oxide surfaces
M. Chen, P.T. Elliott and J.E. Glass (Fargo, ND, USA) 457
Effects of alkyl chain length on the solid state phase behaviors of alkyldimethylamine oxide hydrochlorides and hemihydrochlorides
M. Miyahara, H. Kawasaki, T. Fukuda (Fukuoka, Japan), Y. Ozaki (Nishinomiya, Japan) and H. Maeda (Fukuoka, Japan) 475
Alkanediyl-α,ω-bis(dimethylalkylammonium bromide) surfactants: Part 8. Pressure-jump study of the kinetics of micellar equilibria in aqueous solutions of alkanediyl-α,ω-bis(dimethyldodecylammonium bromide) surfactants
W. Ulbricht (Bayreuth, Germany) and R. Zana (Strasbourg, France) 487
Bilayer fragments and bilayered micelles (bicelles) of dimyristoylphosphatidylglycerol (DMPG) are induced by storage in distilled water at 4°C
H.W. Meyer, W. Richter (Jena, Germany), W. Rettig (Halle, Germany) and M. Stumpf (Jena, Germany) 495
Bending deformation of neutral polymer gels induced by electric fields
J. Fehér, G. Filipcsei, J. Szalma and M. Zrínyi (Budapest, Hungary) 505
High-frequency shear and volume viscoelastic moduli of casein particle gel
C. Smyth, E.D. Kudryashov and V. Buckin (Dublin, Ireland) 517
Viscous and elastic properties of a cationic and a zwitterionic drag reducing surfactant
P. Štern and J. Myška (Praha, Czech Republic) 527
Phase changes during silica particle formation in water-in-oil emulsions
J. Esquena and C. Solans (Barcelona, Spain) 533
Lecithin organogel: A micellar system with unique properties
Y.A. Shchipunov (Vladivostok, Russia) 541

Aggregates of sodium hyaluronate with cationic and aminoxide surfactants in aqueous solution — light scattering study
M. Pisárčik (Nagoya, Japan; Bratislava, Slovakia), T. Imae (Nagoya, Japan), F. Devínsky and I. Lacko (Bratislava, Slovakia) 555
Influence of sodium dodecyl sulfate on structure and rheology of aqueous solutions of the nonionic surfactant tetraethyleneglycol-monododecyl ether ($C_{12}E_4$)
F. Nettesheim, J. Zipfel (Freiburg i. Br., Germany), P. Lindner (Grenoble, France) and W. Richtering (Kiel, Germany) 563
Determination of the surface area of silver sols based on the rate of the electrocatalytic oxidation of *N*,*N*-dimethyl-*p*-phenylenediamine with chloropentaamine-cobalt(III) chloride
U. Nickel, C. Liu and Y. Chen (Erlangen, Germany) 575
Amphiphilic alkyl dimethylaminopyridinium compounds — on the design of SHG active cationic amphiphiles and their adsorption behavior: Part 1. Synthesis of SHG-active alkyl dimethylaminopyridinium bromide, structure and physical properties
K. Haage, H. Motschmann, S.-M. Bae (Potsdam, Germany) and E. Gründemann (Berlin-Adlershof, Germany) 583
Rheological and optical properties in systems containing TiO_2 nanoparticles
S.D. Romano, E.O. Acosta (Buenos Aires, Argentina), T. Dürrschmidt (Bayreuth, Germany) and D.H. Kurlat (Buenos Aires, Argentina) 595
Adsorption layer structures and spreading behavior of aqueous non-ionic surfactants on graphite
T. Svitova (Berkeley, CA, USA; Moscow, Russia), R.M. Hill (Midland, MI, USA) and C.J. Radke (Berkeley, CA, USA) 607
Peculiarities in the physico-chemical behaviour of non-statistically substituted carboxymethylcelluloses
J. Kötz, I. Bogen (Potsdam, Germany), T. Heinze, U. Heinze (Jena, Germany), W.-M. Kulicke and S. Lange (Hamburg, Germany) 621
Thermodynamic modeling of precipitation behavior in mixed micellar systems
N.A. Smirnova (St. Petersburg, Russia) 635
Solubilization of alcohols in SDS and TTAB from isentropic partial molar compressibilities and solubilities
H. Høiland, M.I. Gjerde, C. Mo and E. Lie (Bergen, Norway). 651
Experimental and theoretical studies of the system n-decyl-β-D-maltopyranoside + water
H. Kahl, S. Enders and K. Quitzsch (Leipzig, Germany) 661
Phase inversion of W/O emulsions by adding hydrophilic surfactant — a technique for making cosmetics products
D.J. Miller, T. Henning and W. Grünbein (Frankfurt am Main, Germany) 681
Interactions of tetradecyldimethylaminoxide with polyacrylic and polymethacrylic acids in aqueous solution
F.P. Cavasino (Palermo, Italy), H. Hoffmann (Bayreuth, Germany), C. Sbriziolo (Palermo, Italy) and M.L. Turco Liveri (Palermo, Italy; Bayreuth, Germany) 689
The influence of water on phase transitions of a drug-loaded reverse micellar solution into lamellar liquid crystals
S. Mackeben, M. Müller and C.C. Müller-Goymann (Braunschweig, Germany) 699
Structural properties of cationic surfactant/pentanol/water systems and their interaction with negatively charged layer silicates
R. Patakfalvi and I. Dékány (Szeged, Hungary) 715
Control of particle morphology and film structures of carboxylated poly (*n* butylacrylate)/poly (methyl methacrylate) composite latex particles
S. Kirsch, A. Pfau (Ludwigshafen, Germany), J. Stubbs and D. Sundberg (Durham, NH, USA) 725
Physicochemical properties of resorcinol-type calix[4]arenes phosphoric esters having four alkyl side chains in aqueous solution
K. Esumi, K. Shoji, K. Torigoe (Tokyo, Japan), Y. Koide and H. Shosenji (Kumamoto, Japan) 739
Effects of fluoroalkyl chain length and added moles of oxyethylene on aggregate formation of branched-tail fluorinated anionic surfactants
M. Sagisaka, A. Ito (Chiba, Japan), Y. Kondo, N. Yoshino (Tokyo, Japan), K. Ok Kwon (Inchon City, South Korea), H. Sakai and M. Abe (Chiba, Tokyo, Japan) 749
Electrochemical control of solubilization using a ferrocene-modified nonionic surfactant
T. Takei (Chiba, Japan), H. Sakai (Chiba, Tokyo, Japan), Y. Kondo, N. Yoshino (Tokyo, Japan) and M. Abe (Chiba, Tokyo, Japan) 757
Nitroxide spin probes for magnetic resonance characterization of ordered systems
I. Dragutan, V. Dragutan, A. Caragheorgheopol (Bucharest, Romania), A.K. Zarkadis (Ioannina, Greece), H. Fischer (Konstanz, Germany) and H. Hoffmann (Bayreuth, Germany) 767
Rheological properties of acid gel cleansers
M. Pflaumbaum, F. Müller, J. Peggau, W. Goertz and B. Grüning (Essen, Germany) 777
Influence of polymeric surfactants on pesticidal suspension concentrates: dispersing ability, milling efficiency and stabilization power
S. Haas, H.-W. Hässlin (Münchwilen, CH, USA) and C. Schlatter (Greensboro, USA). 785

Multianalytical approach to characterise adsorption of industrial grade surfactants or polymers on different crystalline agrochemicals
M. von Raumer (Basel, Switzerland), H.-W. Haesslin (Muenchwilen, Switzerland) and R. Hilfiker (Basel, Switzerland) 795
The impact of rheological modifiers on water-borne coatings
U. Kästner (Memmingen, Germany) 805

Calendar 823
Author Index. 825
Subject Index. 827
Volume Contents 831
Contents Continued 835